

Nontrivial RR two-form field strength and SU(3)-structure

Peter Kaste, Ruben Minasian, Michela Petrini and Alessandro Tomasiello

Centre de Physique Théorique, Ecole Polytechnique¹
91128 Palaiseau Cedex, France

kaste, ruben, petrini, tomasiel@cpht.polytechnique.fr

Abstract: We discuss how in the presence of a nontrivial RR two-form field strength and nontrivial dilaton the conditions of preserving supersymmetry on six-dimensional manifolds lead to generalized monopole and Killing spinor equations. We show that the manifold is Kähler in the ten-dimensional string frame if $F_0^{(1,1)} = 0$. We then determine explicitly the intrinsic torsion of the SU(3)-structure on six-manifolds that result via Kaluza-Klein reduction from seven-manifolds with G_2 -structure of generic intrinsic torsion. Lastly we give explicitly the intrinsic torsion of the SU(3)-structure for an $\mathcal{N}=1$ supersymmetric background in the presence of nontrivial RR two-form field strength and nontrivial dilaton.

1 Introduction

A better understanding of $\mathcal{N}=1$ supersymmetric compactifications of string theory to four dimensions is an important step towards more realistic string theories. A promising avenue to take is to break the $\mathcal{N}=2$ supersymmetry of the well-studied Calabi-Yau compactifications of type II string theories down to $\mathcal{N}=1$ by including a background of RR field strength that may describe either internal RR fluxes or spacetime filling D-branes. Of course their presence back-reacts on the metric. In particular the new supersymmetric ground state is no longer a Calabi-Yau manifold. It is an interesting question to study the geometry of these minimally supersymmetric ground states and to characterize how their structure deviates from the one of a Calabi-Yau.

In [1] these questions have been addressed for compactifications of type IIA with a background of nontrivial RR two-form field strength and nontrivial dilaton. This is the situation that is easiest to analyze, since the triple (g, F, φ) of the groundstate metric g , the background two-form field strength F and the dilaton φ can be described in terms of a G_2 manifold Y . Namely this is the internal manifold of the purely geometrical M-theory compactification which via Kaluza-Klein reduction gives rise to the above type IIA configuration.

¹Unité mixte du CNRS et de l'EP, UMR 7644

In these proceedings we first construct explicitly an $SU(3)$ -structure (g_X, J, ψ_3) on the six-dimensional base space X obtained by Kaluza-Klein reduction from a seven-dimensional manifold Y with G_2 -structure. We then analyze the constraints that preserving $\mathcal{N}=1$ supersymmetry in four dimensions imposes on this $SU(3)$ -structure by requiring the G_2 -structure on Y to be torsion-free. These constraints can be cast into a primitivity constraint on F , a monopole equation relating $d\varphi$ to F and a Killing spinor equation on the associated $SU(3)$ -invariant spinor. It immediately follows that the manifold (X, g_X, J) is Kähler in the string frame if the primitive part of $F^{(1,1)}$ vanishes.

Next we compute explicitly the intrinsic torsion of the $SU(3)$ -structure (g_X, J, ψ_3) on X for a generic G_2 -structure on Y . Specializing this result to the case of torsion-free G_2 -structure, we give the intrinsic torsion of the $SU(3)$ -structure for an $\mathcal{N}=1$ supersymmetric background in the presence of nontrivial RR two-form field strength and nontrivial dilaton. Concretely, we show that in the notation of [2] its components are given by $W_1 = W_2^- = W_3 = 0$, $W_2^+ = -\check{F}_0^{(1,1)}$, $W_4 = -(\beta - 2\alpha)d\varphi$ and $W_5 = -(\beta - 3\alpha)d\varphi$.

This intrinsic torsion is the obstruction for the Levi-Civita connection of (X, g_X) to have holonomy $SU(3)$. It can therefore be seen as a measure of how the manifold fails to be a Calabi-Yau. Recently the concept of intrinsic torsion of G -structures has been applied to compactifications with background fields in [3, 4, 5, 6, 7]. In [4] it is in particular argued to describe the mirror of NS three-form fluxes in an otherwise purely geometrical mirror compactification. For $\mathcal{N}=1$ supersymmetric compactifications to three dimensions in the presence of background fields F and φ and for further references we refer to [1].

2 From a G_2 -structure to an $SU(3)$ -structure

Let Y be the seven-dimensional manifold to which one lifts in M-theory and g_Y the background metric on Y . Via Kaluza-Klein reduction it is related to the background metric g_X on the internal space X of the type IIA compactification by

$$ds_Y^2 = e^{-2\alpha\varphi} ds_X^2 + e^{2\beta\varphi} (dz + A)^2, \quad (2.1)$$

where A is the RR one-form potential and φ the dilaton of type IIA. The parameters α and β determine the frame of g_X in ten-dimensional type IIA. They take the values $(\alpha, \beta) = (1/3, 2/3)$ for the string frame.

Suppose Y carries a G_2 -structure. The latter is specified by the doublet (g_Y, Φ) where Φ is a G_2 -invariant, nowhere vanishing three-form on Y . It can be represented as

$$\Phi = \frac{1}{3!} \phi_{ABC} \hat{e}^A \hat{e}^B \hat{e}^C, \quad (2.2)$$

where $\hat{e}^A$ for $A = 1, \dots, 7$ is a frame of orthonormal one-forms w.r.t. g_Y and ϕ_{ABC} are the structure constants of the imaginary octonions. Moreover, the G_2 -structure singles out a unique G_2 -invariant spinor ϵ . It is real (Majorana) and satisfies

$$\gamma_{AB} \epsilon = i \phi_{ABC} \gamma^C \epsilon \quad \text{and} \quad \phi_{ABC} = -i \epsilon^\dagger \gamma_{ABC} \epsilon. \quad (2.3)$$

If and only if the G_2 -structure is torsion-free, the Levi-Civita connection associated to g_Y has holonomy in G_2 and in that case the spinor ϵ will be covariantly constant w.r.t. the Levi-Civita connection. This spinor ϵ on Y is then the internal part of the supersymmetry

generator in the $\mathcal{N}=1$ supersymmetric M-theory compactification on Y . It is projectible onto X along the $U(1)$ -fibers of the Kaluza-Klein bundle $\pi : Y \mapsto X$ if it is constant along these fibers. In that case it becomes the internal part of the supersymmetry generator in type IIA. Note that since this constant along the fiber may vary over X , the spinor ϵ as seen on X has a $U(1)$ gauge symmetry. In particular X need only carry a Spin^c -structure and not a Spin-structure.

If ϵ is projectible and whether the G_2 -structure is torsion-free or not, the identity (2.3) projects on X onto

$$\gamma_{ab}\epsilon = i\psi_{abc}\gamma^c\epsilon + iJ_{ab}\gamma\epsilon \quad \text{and} \quad \gamma_a\gamma\epsilon = -iJ_{ab}\gamma^b\epsilon, \quad (2.4)$$

where we have defined $\psi_{abc} \equiv \phi_{abc}$ and $J_{ab} \equiv \phi_{ab7}$ for $a, b, c = 1, \dots, 6$ and where $\gamma = \gamma^7$ is the chirality operator on X . Furthermore

$$\psi_{abc} = -i\epsilon^\dagger\gamma_{abc}\epsilon \quad \text{and} \quad J_{ab} = -i\epsilon^\dagger\gamma_{ab}\gamma\epsilon. \quad (2.5)$$

The spinor ϵ on X can hence be used to build nowhere vanishing forms

$$\psi_3 \equiv \frac{1}{3!}\psi_{abc}e^ae^be^c \quad \text{and} \quad J \equiv \frac{1}{2}J_{ab}e^ae^b \quad (2.6)$$

on X . Moreover, since $J_a{}^b J_b{}^c = -\delta_a{}^c$, this $J_a{}^b$ defines a natural almost complex structure on X with respect to which g_X is automatically hermitian, with associated two-form J . Splitting each tangent plane into a holomorphic and antiholomorphic space w.r.t. $J_a{}^b$ one sees that ψ_3 is the real part of a $(3, 0)$ -form $\Omega = \psi_3 - i(*\psi_3)$. In particular the forms ψ_3 and J are not only nowhere vanishing on X , they are also by construction invariant under the action of $SU(3)$ on the tangent bundle TX . Altogether, the triple (g_X, J, ψ_3) defines an $SU(3)$ -structure on X . Its associated $SU(3)$ invariant spinor is ϵ .

In summary, if Y is a (Kaluza-Klein) $U(1)$ -bundle over X and has a G_2 -structure with an associated projectible G_2 -invariant spinor, then the base X carries an $SU(3)$ -structure. This implies that the structure group of TX is $SU(3)$. However, the $SU(3)$ -structure on X will generically have torsion, even if the G_2 -structure was torsion-free. I.e. although there exist connections on TX that are compatible with the metric and have holonomy $SU(3)$, generically none of them will be torsion-free. In that case the Levi-Civita connection cannot have holonomy $SU(3)$ and X is not a Calabi-Yau. The intrinsic torsion of the $SU(3)$ -structure (X, g_X, J, ψ_3) is the obstruction for it to be a Calabi-Yau.

In order to see how X fails to be a Calabi-Yau, we look at the differential equations satisfied by ϵ on X or alternatively by ψ_3 and J .

3 The monopole and Killing spinor equations

Let the G_2 -structure on Y be torsion-free so that we preserve $\mathcal{N}=1$ supersymmetry in four dimensions. The covariant constancy of ϵ on Y reduces to the following system on X

$$\left(D_a + \frac{i}{2}\alpha(\partial_b\varphi)J^b{}_a\gamma\right)\epsilon + i\left(\frac{1}{2}\alpha(\partial_b\varphi)\psi^b{}_{ac} - \frac{1}{4}\tilde{F}_{ab}J^b{}_c\right)\gamma^c\epsilon = 0, \quad (3.1a)$$

$$\left(\frac{1}{4}\tilde{F}_{ab}J^{ab}\right)\gamma\epsilon + \left(\frac{1}{4}\tilde{F}_{ab}\psi^{ab}{}_c - \beta(\partial_a\varphi)J^a{}_c\right)\gamma^c\epsilon = 0, \quad (3.1b)$$

where we have defined $\check{F} \equiv e^{(\alpha+\beta)\varphi} F$ and where D_a denotes the covariant derivative w.r.t. the Levi-Civita connection on (X, g_X) . Since $\gamma^A \epsilon$ are linearly independent, the latter of these equations gives

$$F^{ab} J_{ab} = 0 \quad \Leftrightarrow \quad J \lrcorner F = 0, \quad (3.2a)$$

$$\beta (\partial_a \varphi) J^a_c = \frac{1}{4} \check{F}^{ab} \psi_{abc} \quad \Leftrightarrow \quad \beta d\varphi = \frac{1}{2} \check{F} \lrcorner (*\psi_3), \quad (3.2b)$$

where $\lrcorner$ denotes the contraction of forms w.r.t. the metric g_X . On a Kähler manifold the first of these would imply that F is primitive and even though (X, g_X, J) might not be Kähler we will refer to (3.2a) as a primitivity constraint. The second equation is a generalized monopole equation. It relates $d\varphi$ to the $(2, 0)$ and $(0, 2)$ parts of F w.r.t. the almost complex structure. Inserting these into (3.1a) leads to the following Killing spinor equation on X for $\beta = 2\alpha$,

$$\left(D_a + \frac{i}{2} \alpha (\partial_b \varphi) J^b_a \gamma - \frac{i}{8} [\check{F}_{ab} J^b_c + \check{F}_{cb} J^b_a] \gamma^c \right) \epsilon = 0. \quad (3.3)$$

This implies that the only nonvanishing components of the Nijenhuis tensor associated to the almost complex structure $J_a^b = -i\epsilon^\dagger \gamma_a^b \gamma \epsilon$ are

$$N_{bc}^{\bar{a}} = \frac{i}{2} \left(\check{F}_{cd} \epsilon^{\bar{d}}_{\bar{b}} - \check{F}_{bd} \epsilon^{\bar{d}}_{\bar{c}} \right) \quad \text{and} \quad N_{\bar{b}\bar{c}}^a = -\frac{i}{2} \left(\check{F}_{cd} \epsilon^d_{\bar{b}} - \check{F}_{bd} \epsilon^d_{\bar{c}} \right), \quad (3.4)$$

where we have used the holomorphic/antiholomorphic basis of TX w.r.t. J_a^b . The almost complex structure defined by the spinor ϵ is therefore integrable if and only if $F^{(1,1)} \equiv 0$. Since $dJ = 0$ we find that in this case (X, g_X, J) is Kähler for $\beta = 2\alpha$, i.e. defines a torsion-free $U(3)$ -structure. The Killing spinor equation on X then reduces to

$$\left(D_a + \frac{i}{2} \alpha (\partial_b \varphi) J^b_a \gamma \right) \epsilon = 0. \quad (3.5)$$

Since ϵ is not covariantly constant w.r.t. the Levi-Civita connection for nontrivial φ , $F^{(2,0)}$ and $F^{(0,2)}$, the $SU(3)$ -structure (X, g_X, J, ψ_3) however still has torsion.

4 The general relation between the intrinsic G_2 -torsion and $SU(3)$ -torsion

The intrinsic torsion of the G_2 -structure (g_Y, Φ) takes values in $(\mathfrak{g}_2)_\perp \otimes T^*Y$, where $\mathfrak{so}(7) = \mathfrak{g}_2 \oplus (\mathfrak{g}_2)_\perp$ and can be decomposed as [2]

$$X_1 \in Y \otimes \mathbb{R}, \quad X_2 \in \Lambda_{14}^2 T^*Y, \quad X_3 \in \Lambda_{27}^4 T^*Y, \quad X_4 \in \Lambda_7 T^*Y, \quad (4.1)$$

where $\Lambda_m^n T^*Y$ denotes n -forms that transform in the representation $\underline{m}$ of G_2 . In particular these representations imply that

$$X_2 \lrcorner_{g_Y} \Phi = 0, \quad \Phi \lrcorner_{g_Y} X_3 = 0, \quad (*\Phi) \lrcorner_{g_Y} X_3 = 0, \quad (4.2)$$

where $\lrcorner_{g_Y}$ denotes the contraction of forms w.r.t. the metric g_Y . The components (4.1) are determined through $d\Phi$ and $d(*\Phi)$ as [2]

$$d\Phi = X_1(*\Phi) + X_4 \wedge \Phi + X_3, \quad (4.3a)$$

$$d(*\Phi) = \frac{4}{3}X_4 \wedge (*\Phi) + X_2 \wedge \Phi. \quad (4.3b)$$

Let us furthermore introduce the notation

$$X_j = Y_j + Z_j \wedge \hat{e}^7, \quad \text{for } j = 1, \dots, 4. \quad (4.4)$$

Then the constraints (4.2) take the form

$$X_2 \lrcorner_{g_Y} \Phi = 0 \quad \Leftrightarrow \quad \begin{cases} e^{\alpha\varphi} Y_2 \lrcorner_{g_X} \psi_3 + Z_2 \lrcorner_{g_X} J = 0, \\ J \lrcorner_{g_X} Y_2 = 0, \end{cases} \quad (4.5a)$$

$$\Phi \lrcorner_{g_Y} X_3 = 0 \quad \Leftrightarrow \quad \begin{cases} e^{\alpha\varphi} \psi_3 \lrcorner_{g_X} Y_3 - J \lrcorner_{g_X} Z_3 = 0, \\ \psi_3 \lrcorner_{g_X} Z_3 = 0, \end{cases} \quad (4.5b)$$

$$(*\Phi) \lrcorner_{g_Y} X_3 = 0 \quad \Leftrightarrow \quad \frac{1}{2} e^{\alpha\varphi} J^2 \lrcorner_{g_X} Y_3 + (*\psi_3) \lrcorner_{g_X} Z_3 = 0. \quad (4.5c)$$

In the following all the contractions will be taken w.r.t. the metric g_X and we will drop the label, i.e. $\lrcorner \equiv \lrcorner_{g_X}$.

Analogously the intrinsic torsion of the $SU(3)$ -structure (g_X, J, ψ_3) takes values in $\mathfrak{su}(3)_\perp \otimes T^*X$, where $\mathfrak{so}(6) = \mathfrak{su}(3) \oplus \mathfrak{su}(3)_\perp$. It can be decomposed as

$$\begin{aligned} W_1 &\in X \otimes \mathbb{C}, & W_2 &\in \Lambda_{8 \oplus 8}^{(1,1)} T^*X|_{\text{primitive}}, \\ W_3 &\in (\Lambda_6^{(2,1)} T^*X \oplus \Lambda_6^{(1,2)} T^*X)|_{\text{primitive}}, & W_4 &\in \Lambda_3^{(1,0)} T^*X \oplus \Lambda_3^{(0,1)} T^*X, \\ W_5 &\in \Lambda_3^{(1,0)} T^*X \oplus \Lambda_3^{(0,1)} T^*X, \end{aligned} \quad (4.6)$$

where $\Lambda_m^{(n_1, n_2)} T^*X$ denotes (n_1, n_2) -forms that transform in the representation $\underline{m}$ of $SU(3)$. In particular these representations imply that

$$J \lrcorner W_2 = 0, \quad J \lrcorner W_3 = 0, \quad \psi_3 \lrcorner W_3 = 0 \quad \text{and} \quad (*\psi_3) \lrcorner W_3 = 0. \quad (4.7)$$

We can similarly express the components of its intrinsic torsion through dJ , $d\psi_3$ and $d(*\psi_3)$ as [2]

$$dJ = \frac{3}{2} \text{Im}(\overline{W_1} \Omega) + W_4 \wedge J + W_3 \quad (4.8a)$$

$$= \frac{3}{2} W_1^- \psi_3 - \frac{3}{2} W_1^+ (*\psi_3) + W_4 \wedge J + W_3,$$

$$d\psi_3 = W_1^+ J^2 + W_2^+ \wedge J + \text{Re}(W_5 \wedge \overline{\Omega}) \quad (4.8b)$$

$$= W_1^+ J^2 + W_2^+ \wedge J + W_5^+ \wedge \psi_3 + W_5^- \wedge (*\psi_3),$$

$$d(*\psi_3) = W_1^- J^2 + W_2^- \wedge J + \text{Im}(W_5 \wedge \overline{\Omega}) \quad (4.8c)$$

$$= W_1^- J^2 + W_2^- \wedge J + W_5^+ \wedge (*\psi_3) - W_5^- \wedge \psi_3,$$

$$d\Omega = W_1 J^2 + W_2 \wedge J + \overline{W_5} \wedge \Omega, \quad (4.8d)$$

where we have used the (3,0)-form

$$\Omega = \psi_3 - i(*\psi_3), \quad (4.9)$$

as well as

$$W_j = W_j^+ - iW_j^-, \quad \text{for } j = 1, 2, 5. \quad (4.10)$$

The minus signs on the imaginary parts result from the fact that compared to [2] we use the opposite orientation, where $J = e^{14} + e^{25} + e^{36}$.

Using

$$\Phi = e^{-3\alpha\varphi}\psi + e^{-2\alpha\varphi}J \wedge \hat{e}^7, \quad (4.11a)$$

$$(*\Phi) = e^{-4\alpha\varphi}(*J) + e^{-3\alpha\varphi}(*\psi_3) \wedge \hat{e}^7 = -\frac{1}{2}e^{-4\alpha\varphi}J^2 + e^{-3\alpha\varphi}(*\psi_3) \wedge \hat{e}^7, \quad (4.11b)$$

equations (4.3a), (4.3b) and (4.8a)–(4.8d), as well as $\check{F} = e^{(\alpha+\beta)\varphi}F$, one derives the following four identities,

$$\begin{aligned} & 3\alpha \psi_3 \wedge d\varphi + W_1^+ J^2 + W_2^+ \wedge J + \text{Re}(W_5 \wedge \bar{\Omega}) + J \wedge \check{F} \\ &= -\frac{1}{2}e^{\alpha\varphi}X_1 J^2 - \psi_3 \wedge Y_4 + e^{3\alpha\varphi}Y_3, \end{aligned} \quad (4.12a)$$

$$\begin{aligned} & (\beta - 2\alpha)J \wedge d\varphi + \frac{3}{2}W_1^- \psi_3 - \frac{3}{2}W_1^+(*\psi_3) + W_4 \wedge J + W_3 \\ &= e^{-(\alpha+\beta)\varphi}X_1(*\psi_3) + e^{-\beta\varphi}J \wedge Y_4 - e^{-(\alpha+\beta)\varphi}Z_4\psi_3 + e^{-(\beta-2\alpha)\varphi}Z_3, \end{aligned} \quad (4.12b)$$

$$\begin{aligned} & 2\alpha J^2 \wedge d\varphi - J^2 \wedge W_4 - (*\psi_3) \wedge \check{F} \\ &= -\frac{2}{3}J^2 \wedge Y_4 + e^{\alpha\varphi}\psi_3 \wedge Y_2, \end{aligned} \quad (4.12c)$$

$$\begin{aligned} & (\beta - 3\alpha)d\varphi \wedge (*\psi_3) + W_1^- J^2 + W_2^- \wedge J + \text{Im}(W_5 \wedge \bar{\Omega}) \\ &= -\frac{4}{3}e^{-\beta\varphi}(*\psi_3) \wedge Y_4 - \frac{2}{3}e^{-(\alpha+\beta)\varphi}Z_4 J^2 + e^{(\alpha-\beta)\varphi}J \wedge Y_2 + e^{-\beta\varphi}\psi_3 \wedge Z_2. \end{aligned} \quad (4.12d)$$

From these equations we can project onto the various W_j 's by suitable contractions and express them through the intrinsic torsion (Y_j, Z_j) of the G_2 -structure. To this end it is convenient to decompose forms as

$$F = F^{(0)}J + F_0^{(1,1)} + F^{(2,0)} + F^{(0,2)} \quad (4.13a)$$

$$= \frac{1}{3}(J \lrcorner F)J + F_0^{(1,1)} + \frac{1}{8}[(F \lrcorner \bar{\Omega}) \lrcorner \Omega + (F \lrcorner \Omega) \lrcorner \bar{\Omega}],$$

$$Y_2 = Y_2^{(0)}J + Y_{2,0}^{(1,1)} + Y_2^{(2,0)} + Y_2^{(0,2)} \quad (4.13b)$$

$$= \frac{1}{3}(J \lrcorner Y_2)J + Y_{2,0}^{(1,1)} + \frac{1}{8}[(Y_2 \lrcorner \bar{\Omega}) \lrcorner \Omega + (Y_2 \lrcorner \Omega) \lrcorner \bar{\Omega}],$$

$$Z_3 = Z_3^{(0)}\psi_3 + \tilde{Z}_3^{(0)}(*\psi_3) + (Z_3^{(1,0)} + Z_3^{(0,1)}) \wedge J + Z_{3,0}^{(2,1)} + Z_{3,0}^{(1,2)} \quad (4.13c)$$

$$= \frac{1}{4}(\psi_3 \lrcorner Z_3)\psi_3 + \frac{1}{4}((*\psi_3) \lrcorner Z_3)(* \psi_3) + \frac{1}{2}(J \lrcorner Z_3) \wedge J + Z_{3,0}^{(2,1)} + Z_{3,0}^{(1,2)},$$

$$Y_3 = Y_3^{(0,1)} \wedge \Omega + Y_3^{(1,0)} \wedge \bar{\Omega} + (Y_3^{(2,0)} + Y_3^{(0,2)}) \wedge J + Y_{3,0}^{(2,2)} + Y_{3,0}^{(1,1)} \wedge J + Y_3^{(0)}J^2, \quad (4.13d)$$

where subscripts “0” denote primitive forms. Using projectors such as

$$Y_{3,0}^{(1,1)} = J \lrcorner Y_3 - \frac{1}{3}(J^2 \lrcorner Y_3)J - \frac{i}{8}(\Omega \lrcorner Y_3) \lrcorner \bar{\Omega} + \frac{i}{8}(\bar{\Omega} \lrcorner Y_3) \lrcorner \Omega, \quad (4.14)$$

the components W_j of the intrinsic torsion of the $SU(3)$ -structure (g_X, J, ψ_3) can be expressed through the components (Y_j, Z_j) of the intrinsic torsion of the G_2 -structure (g_Y, Φ) as,

$$W_1^+ = -\frac{2}{3}e^{-(\alpha+\beta)\varphi}X_1 - \frac{2}{3}e^{-(\beta-2\alpha)\varphi}\tilde{Z}_3^{(0)}, \quad (4.15a)$$

$$W_1^+ + \check{F}^{(0)} = -\frac{1}{2}e^{\alpha\varphi}X_1 + e^{3\alpha\varphi}Y_3^{(0)}, \quad (4.15b)$$

$$W_1^- = -\frac{2}{3}e^{-(\alpha+\beta)\varphi}Z_4, \quad (4.15c)$$

$$W_2^+ + \check{F}_0^{(1,1)} = e^{3\alpha\varphi}Y_{3,0}^{(1,1)}, \quad (4.15d)$$

$$W_2^- = e^{(\alpha-\beta)\varphi}Y_{2,0}^{(1,1)}, \quad (4.15e)$$

$$W_3 = e^{-(\beta-2\alpha)\varphi}[Z_{3,0}^{(2,1)} + Z_{3,0}^{(1,2)}], \quad (4.15f)$$

$$W_4 + (\beta - 2\alpha)d\varphi = e^{-\beta\varphi}Y_4 + e^{-(\beta-2\alpha)\varphi}[Z_3^{(1,0)} + Z_3^{(0,1)}], \quad (4.15g)$$

$$W_4 - 2\alpha d\varphi + \frac{1}{2}\check{F} \lrcorner (*\psi_3) = \frac{2}{3}Y_4 - \frac{1}{2}e^{\alpha\varphi}Y_2 \lrcorner \psi_3, \quad (4.15h)$$

$$W_5^{(1,0)} - 3\alpha d\varphi^{(1,0)} + \frac{i}{4}\check{F} \lrcorner \Omega = Y_4^{(1,0)} - \frac{1}{4}e^{3\alpha\varphi}\Omega \lrcorner Y_3, \quad (4.15i)$$

$$W_5^{(1,0)} + (\beta - 3\alpha)d\varphi^{(1,0)} = \frac{4}{3}e^{-\beta\varphi}Y_4^{(1,0)} - \frac{1}{4}e^{(\alpha-\beta)\varphi}Y_2 \lrcorner \Omega - ie^{-\beta\varphi}Z_2^{(1,0)}. \quad (4.15j)$$

The decomposition of the X_j 's into the W_j 's at the level of representations figures already in [2]. Equations (4.15a)–(4.15j) determine the explicit coefficients that appear in this decomposition for a metric (2.1) of the form that appears in Kaluza-Klein reductions to an arbitrary frame.

Let's now specialize again to the case of torsion-free G_2 -structure, where all the right hand sides vanish. From (4.15a) and (4.15b) we recover the primitivity constraint (3.2a)

$$\frac{1}{3}J \lrcorner F \equiv F^{(0)} = 0, \quad (4.16)$$

whereas equations (4.15g), (4.15h) as well as (4.15i),(4.15j) and their complex conjugates give us the monopole equation (3.2b)

$$\beta d\varphi = \frac{1}{2}\check{F} \lrcorner (*\psi_3). \quad (4.17)$$

In addition to these two constraints, the components of the intrinsic torsion of the $SU(3)$ -structure in the case of torsion-free G_2 -structure are given by

$$\begin{aligned} W_1^\pm &= 0, & W_2^- &= 0, & W_3 &= 0, \\ W_2^+ &= -\check{F}_0^{(1,1)}, & W_4 &= -(\beta - 2\alpha)d\varphi, & W_5 &= -(\beta - 3\alpha)d\varphi. \end{aligned} \quad (4.18)$$

Moreover, we recover the result from the previous section that for $\beta = 2\alpha$ the manifold (X, g_X, J) is Kähler if $F_0^{(1,1)} = 0$, since then the only nonvanishing component of the intrinsic torsion of (g_X, J, ψ_3) is W_5 .

Acknowledgement

We would like to thank B. Acharya, D. Calderbank, M. Douglas, D. Martelli and A. Moroianu for useful discussions. PK would furthermore like to thank the organizers of the 35th Ahrenschoop Symposium for the invitation to participate in a stimulating conference and the opportunity to present this work. This work is supported in part by EU contract HPRN-CT-2000-00122 and by INTAS contracts 55-1-590 and 00-0334. PK and MP are supported by European Commission Marie Curie Postdoctoral Fellowships under contract numbers HPMF-CT-2000-00919 and HPMF-CT-2001-01277.

References

- [1] P. Kaste, R. Minasian, M. Petrini and A. Tomasiello, “Kaluza-Klein bundles and manifolds of exceptional holonomy,” JHEP **0209** (2002) 033, arXiv:hep-th/0206213.
- [2] S. Chiossi, S. Salamon, “The intrinsic torsion of $SU(3)$ and G_2 structures” Proc. conf. Differential Geometry Valencia 2001, math.DG/0202282.
- [3] J.P. Gauntlett, D. Martelli, S. Pakis and D. Waldram, “G-structures and wrapped NS5-branes,” arXiv:hep-th/0205050.
- [4] S. Gurrieri, J. Louis, A. Micu and D. Waldram, “Mirror symmetry in generalized Calabi-Yau compactifications,” arXiv:hep-th/0211102.
- [5] G.L. Cardoso, G. Curio, G. Dall’Agata, D. Lüst, P. Manousselis, G. Zoupanos, “Non-Kähler string backgrounds and their five torsion classes,” arXiv:hep-th/0211118.
- [6] J.P. Gauntlett and S. Pakis, “The geometry of d=11 Killing spinors,” arXiv:hep-th/0212008.
- [7] E. Goldstein and S. Prokushkin, “Geometric model for complex non-Kähler manifolds with $SU(3)$ -structure,” arXiv:hep-th/0212307.
- [8] M. Cvetič, G. W. Gibbons, H. Lu and C. N. Pope, “Almost special holonomy in type IIA and M theory,” Nucl. Phys. **B 638** (2002) 186, arXiv:hep-th/0203060.